

is imodium a prescription drug

[\[PDF\] how much does bactrim cost at cvs](#)

[\[PDF\] where to buy metronidazole pills](#)

[\[PDF\] tramadol generic 50mg](#)

[\[PDF\] cost omnicef walmart](#)

[\[PDF\] where can i buy liquid tadalafil](#)

[\[PDF\] is cefdinir a generic drug](#)

[\[PDF\] what mg is prescription ibuprofen](#)

Take Imodium with a full glass of water. You should not use Imodium if you have ulcerative colitis, bloody or tarry stools, diarrhea with a high fever, or diarrhea caused by antibiotic medication. Avoid driving or hazardous activity until you know how this medicine will affect you. The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Everyone has diarrhea from time to time, while some people suffer more frequently. Be kind to your digestive system. Treat diarrhea early to get back to your day. To view content sources and attributions, please refer to our editorial policy. Store at room temperature away from moisture and heat. Do not give Imodium to a child younger than 2 years old. Imodium loperamide slows the rhythm of digestion so that the small intestines have more time to absorb fluid and nutrients from the foods you eat. Imodium is also used to reduce the amount of stool in people who have an ileostomy re-routing of the bowel through a surgical opening in the stomach. Your reactions could be impaired. We can all get struck by stomach flu when traveling. Learn about Imodium (Loperamide Hcl) may treat, uses, dosage, side effects, drug interactions, warnings, patient labeling, reviews, and related medications. Children: In children 2 to 5 years of age (20 kg or less), the non-prescription liquid formulation (IMODIUM (loperamide hcl) A-D 1 mg/5 mL) should be used; for. You can buy over-the-counter (OTC) medicines without a prescription from your doctor. Some OTC loperamide (1 brand name: Imodium); bismuth subsalicylate (2 brand names: Kaopectate, Pepto-Bismol). Bismuth Before you take an OTC antidiarrheal medicine, read the directions on the drug facts label. This will tell. Children: In children 2 to 5 years of age (20 kg or less), the non-prescription liquid formulation (IMODIUM A-D 1 mg/ mL) should be used; for ages 6 to 12, either IMODIUM Capsules or IMODIUM A-D Liquid may be used. For children 2 to 12 years of age, the following schedule for capsules or liquid will usually fulfill. 5 days ago - Important Information. You should not use Imodium if you have ulcerative colitis, bloody or tarry stools, diarrhea with a high fever, or diarrhea caused by antibiotic medication. Do not take more than the recommended dose of this medicine. Misuse of loperamide can cause serious heart problems that could. At-home detox attempts using Imodium or other methods should be avoided. Even though Imodium is an opiate it does not work as a replacement medication. Loperamide (Imodium) is a medication for the treatment and relief of acute, chronic, or traveler's diarrhea. Imodium usually cures diarrhea in a couple of days. Common side effects of Imodium are fatigue, dizziness, abdominal pain, and constipation. If diarrhea lasts longer, contact your doctor. Imodium also has the potential to. Nov 5, - Loperamide is the generic form of the brand-name drug Imodium, a medication used to treat diarrhea. Imodium works by slowing down the movement of the gut. This decreases the number of bowel movements and makes stool less watery. Imodium may also be used to decrease the amount of discharge in. Prescription loperamide and over-the-counter (OTC) loperamide are the same strength but available in different dosage forms. OTC loperamide OTC loperamide comes in three forms: two mg. tablet, oral solution at two mg./10 ml. and What's the Difference Between a Brand-Name Drug and a Generic Drug? What's the. May 10, - The opioid epidemic has led to a form of drug abuse that few experts saw coming: the use of Imodium and other anti-diarrhea medications to get high. As efforts to limit prescription opioids intensify, a handful of experts are concerned that more addicts might turn to loperamide much as an alcoholic. Jump to Are there any other precautions or warnings for this medication? - These factors may affect how you should use this medication. Acquired immune deficiency syndrome (AIDS): People with AIDS should stop taking loperamide and contact their doctor if they experience abdominal swelling or distention.