

prozac pharmacological classification

[\[PDF\] when did generic levaquin become available](#)

[\[PDF\] cost of prozac in south africa](#)

[\[PDF\] buy generic tretinoin gel](#)

[\[PDF\] when is abilify generic](#)

[\[PDF\] what does 1mg generic xanax look like](#)

[\[PDF\] hydrocodone generic vs vicodin](#)

[\[PDF\] generic brand for amoxil](#)

Benzodiazepines such as alprazolam , lithium, tricyclic antidepressants: Do not start or stop taking this medicine during pregnancy without your doctor's advice. You must wait at least 14 days after stopping an MAO inhibitor before you can take Prozac. If you also take olanzapine Zyprexa , read the Zyprexa medication guide and all patient warnings and instructions provided with that medication. Not all possible interactions are listed here. This includes prescription and over-the-counter medicines, vitamins, and herbal products. You must wait 5 weeks after stopping fluoxetine before you can take thioridazine or an MAOI. Ask your doctor how to safely stop using this medicine. Increases risk of serotonin syndrome. Increases phenytoin levels and risk of toxicity. Metabolized primarily in the liver to active metabolites. If you miss a dose of Prozac Weekly, take the missed dose as soon as you remember and take the next dose 7 days later. May reverse or decrease fluoxetine effects. This is not a complete list of side effects and others may occur. To reduce risk of intentional overdose, give the smallest quantity of pulvules consistent with good management. The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Subscribe to receive email notifications whenever new articles are published. Activated charcoal, which may be used with sorbitol, may be as effective as emesis or lavage.PROZAC (Fluoxetine) drug information & product resources from MPR including dosage information, educational materials, & patient assistance. The long elimination half-lives of fluoxetine and norfluoxetine assure that, even when dosing is stopped, active drug substance will persist in the body for . The following table provides the outcome classification by treatment group on the Clinical Global Impression (CGI) improvement scale for Studies 1 and 2 combined. Apr 3, - Prozac (fluoxetine) is used to treat major depressive disorder, bulimia, obsessive-compulsive disorder and panic disorder. Includes Prozac side effects, interactions and indications. Apr 3, - Fluoxetine is used to treat depression or obsessive-compulsive disorder in adults. Learn about side effects, interactions and indications. Learn about Prozac (Fluoxetine Hcl) may treat, uses, dosage, side effects, drug interactions, warnings, patient labeling, reviews, and related medications. Find patient medical information for Prozac Oral on WebMD including its uses, side effects and safety, interactions, pictures, warnings and user ratings. Jump to Side effects - The FDA requires Prozac to come with a black box warning stating that antidepressants may increase the risk of suicide in people younger than 25 years. It can lead to suicidal thoughts, or a worsening of these, in children and young adults. Other possible side effects include: decreased libido. Aug 15, - Fluoxetine: learn about side effects, dosage, special precautions, and more on MedlinePlus. fluoxetine. Prozac, Prozac Weekly, Sarafem. Pharmacologic classification: selective serotonin reuptake inhibitor. Therapeutic classification: antidepressant. Pregnancy risk category C. Fluoxetine hydrochloride is the first agent of the class of antidepressants known as selective serotonin-reuptake inhibitors (SSRIs). Despite distinct structural differences between compounds in this class, SSRIs possess similar pharmacological activity. As with other antidepressant agents, several weeks of therapy may be.