

# fluoxetine prescription drug

[\[PDF\] viagra generic indonesia](#)

[\[PDF\] purchase indomethacin online](#)

[\[PDF\] buy elocon ointment](#)

[\[PDF\] quanto costa sporanox compresse](#)

[\[PDF\] lansoprazole ratiopharm 15 mg](#)

[\[PDF\] domperidone online canada](#)

[\[PDF\] is generic imitrex as effective](#)

Mayo Clinic does not endorse companies or products. Do not use fluoxetine if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine. Any use of this site constitutes your agreement to the Terms and Conditions and Privacy Policy linked below. Google Chrome Open the browser. Free E-newsletter Subscribe to Housecall Our general interest e-newsletter keeps you up to date on a wide variety of health topics. Taking fluoxetine with other drugs that make you sleepy or slow your breathing can cause dangerous side effects or death. You should not use fluoxetine if you are allergic to it, if you also take pimozide or thioridazine, or if you are being treated with methylene blue injection. It may take up to 4 weeks before your symptoms improve. Taking an SSRI antidepressant during pregnancy may cause serious lung problems or other complications in the baby. Keep using the medication as directed and tell your doctor if your symptoms do not improve. Report any new or worsening symptoms to your doctor, such as: How to upgrade your browser to Transport Layer Security 1. Your doctor should check your progress at regular visits. This medication may improve your mood, sleep, appetite, and energy level and may help restore your interest in daily living. It may decrease fear, anxiety, unwanted thoughts, and the number of panic attacks. It may also reduce the urge to perform repeated tasks (compulsions such as hand-washing, counting, and checking). If fluoxetine is added to the treatment regimen of a patient already receiving a drug metabolized by CYP2D6, the need for decreased dose of the original medication should be considered. Drugs with a narrow therapeutic index represent the greatest concern (e.g., flecainide, propafenone, vinblastine, and TCAs). Due to the. Apr 3, - Fluoxetine is used to treat major depressive disorder, bulimia nervosa (an eating disorder), obsessive-compulsive disorder, panic disorder, and premenstrual dysphoric disorder (PMDD). Fluoxetine is sometimes used together with another medication called olanzapine (Zyprexa) to treat manic depression. Fluoxetine, also known by trade names Prozac and Sarafem, among others, is an antidepressant of the selective serotonin reuptake inhibitor (SSRI) class. It is used for the treatment of major depressive disorder, obsessive compulsive disorder (OCD), bulimia nervosa, panic disorder and premenstrual dysphoric rubeninorchids.comical half-life?: ?13 days (acute); 46 days. Defines the medication fluoxetine (Prozac, Sarafem, Prozac Weekly) a drug used in the treatment of depression, obsessive-compulsive disorders, panic attacks, and premenstrual dysphoric disorder. Article includes descriptions, uses, drug interactions, and side effects.?What is fluoxetine, and how ?What are the side effects of. Medicare coverage and pricing details for Fluoxetine. Learn more about Medicare prescription drug plans and savings with GoodRx. Fluoxetine (Prozac) is an inexpensive drug used to treat mood problems such as depression, obsessive compulsive disorder, and panic attacks. Generic fluoxetine is covered by most Medicare and insurance plans, but some pharmacy coupons or cash prices may be lower. Pay as little as \$28 per prescription. See Tips. Aug 15, - The doctor or pharmacist will give you the manufacturer's patient information sheet (Medication Guide) when you begin treatment with fluoxetine. Read the information carefully and ask your doctor or pharmacist if you have any questions. You also can obtain the Medication Guide from the FDA website. Mar 17, - Stopping suddenly can cause serious side effects. Your doctor will help you safely stop the medication. Don't take Prozac or other forms of fluoxetine if you're allergic to fluoxetine or any of the inactive ingredients in the various formulations. Cases of severe allergic reactions have been reported and could. Capsule. Read the Medication Guide that comes with PROZAC before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk with your healthcare provider if there is.