

fenofibrate pharmacology class

[\[PDF\] benadryl for dogs generic](#)

[\[PDF\] ribavirin price in india](#)

[\[PDF\] price for benadryl](#)

[\[PDF\] viagra thailand online](#)

[\[PDF\] zolpidem tartrate buy online](#)

[\[PDF\] cetirizine hcl cost](#)

[\[PDF\] cheapest xenical uk](#)

Subscribe to receive email notifications whenever new articles are published. Get emergency medical help if you have any signs of an allergic reaction to fenofibrate: Some brands of fenofibrate should be taken with meals to help your body better absorb the medicine. Your liver and gallbladder function may also need to be checked. Use cyclosporine and fenofibrate cautiously together. Browse drugs by section Browse drugs by manufacturer. Take fenofibrate exactly as it was prescribed for you. Fenofibrate may stimulate the breakdown of triglyceride-rich protein. Store at room temperature away from moisture, heat, and light. Stay signed in for 30 days. Start with 67 mg daily. If overdose does occur, initiate supportive measures. No data are available on use of statins with fenofibrate; however, because of risk of myopathy, rhabdomyolysis, and acute renal failure reported with combined use of statins and gemfibrozil another fibrate derivative, determine whether the benefit of further alterations in lipid levels outweighs the risk of adverse effects. You should not use fenofibrate if you are allergic to it, or if you have liver disease, gallbladder disease, severe kidney disease, or if you are on dialysis. Get your prescription refilled before you run out of medicine completely. Call your doctor for medical advice about side effects. Search UniChem for chemicals with the same backbone. Pharmacology. Indication. For use as adjunctive therapy to diet to reduce elevated LDL-C, Total-C, Triglycerides and Apo B, and to increase HDL-C in adult patients with primary hypercholesterolemia or mixed dyslipidemia (Fredrickson Types IIa and IIb). Structured Indications. Hypercholesterolemia, Familial Mixed ?Identification ?Pharmacology ?References ?Economics. TRICOR (Fenofibrate) drug information & product resources from MPR including dosage information, educational materials, & patient assistance. TRIGLIDE (Fenofibrate) drug information & product resources from MPR including dosage information, educational materials, & patient assistance. TRIGLIDE Rx. Add Drug To My List Compare to related Drugs View/edit/Compare drugs in my list Pharmacological Class: Fibrate. Fenofibrate, marketed as Tricor and under several other brand names, is a drug of the fibrate class. It is mainly used to reduce cholesterol levels in people at risk of cardiovascular disease. Like other fibrates, it reduces both low-density lipoprotein (LDL) and very low density lipoprotein (VLDL) levels, as well as increasing ?Medical uses ?Interactions ?Mechanism of action ?Formulations. The mechanism of action of fenofibrate is as a Peroxisome Proliferator-activated Receptor alpha Agonist. The chemical classification of fenofibrate is PPAR alpha. FDA Pharmacology Summary from FDA Pharm Classes. Fenofibrate is a synthetic phenoxy-isobutyric acid derivate and prodrug with antihyperlipidemic activity. Brand Names: Antara, Fenoglide, Lipofen, Lofibra, TriCor, Triglide, Lipidil Micro, Dom-Fenofibrate, Lipidil Supra, Lipidil EZ. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement. Professional guide for Fenofibrate. Includes: pharmacology Fenofibrate. Pronunciation: FEN-oh-FYE-brate. Class: Antihyperlipidemic Elderly. Pharmacokinetic studies indicate that fenofibrate can be used in elderly subjects who have healthy renal function without accumulation of the drug or metabolites. Learn about Tricor (Fenofibrate) may treat, uses, dosage, side effects, drug interactions, warnings, patient labeling, reviews, and related medications. numbers have been observed in humans after treatment with other members of the fibrate class when liver biopsies were compared before and after treatment in the same. to the drug. In rodents, one to six times the maximum human dosage of fenofibrate increased the incidence of hepatic and pancreatic cancers. Monkeys have not demonstrated these effects in preclinical studies after 12 months of fenofibrate treatment. In humans, other members of the fibrate class have produced observed. fenofibrate (micronized) TriCor. Pharmacologic classification: fibric acid derivative. Therapeutic classification: antihyperlipidemic Interactions Drug-drug. Bile acid resins: May bind with and inhibit absorption of fenofibrate. Fenofibrate should be taken 1 hour before or 4 to 6 hours after taking these agents. Coumarin-type.