

olanzapine borderline personality

In general, attention must be paid to adverse effects. To assess the effects of drug treatment in BPD patients. The findings were suggestive in supporting the use of second-generation antipsychotics, mood stabilisers, and omega-3 fatty acids, but require replication, since most effect estimates were based on single studies. A significant decrease in body weight was observed with topiramate treatment. Conclusions have to be drawn carefully in the light of several limitations of the RCT evidence that constrain applicability to everyday clinical settings among others, patients' characteristics and duration of interventions and observation periods. Data indicated better outcomes for phenelzine sulfate but no significant differences in the other comparisons, except olanzapine which showed more weight gain and sedation than fluoxetine. There was a possible increase in self-harming behaviour, significant weight gain, sedation and changes in haemogram parameters with olanzapine. First-generation antipsychotics flupenthixol decanoate, haloperidol, thiothixene ; second-generation antipsychotics aripiprazole, olanzapine, ziprasidone , mood stabilisers carbamazepine, valproate semisodium, lamotrigine, topiramate , antidepressants amitriptyline, fluoxetine, fluvoxamine, phenelzine sulfate, mianserin , and dietary supplementation omega-3 fatty acid were tested. In addition, the studies may not adequately reflect several characteristics of clinical settings among others, patients' characteristics and duration of interventions and observation periods. Data collection and analysis: Direct drug comparisons comprised two first-generation antipsychotics loxapine versus chlorpromazine , first-generation antipsychotic against antidepressant haloperidol versus amitriptyline; haloperidol versus phenelzine sulfate , and second-generation antipsychotic against antidepressant olanzapine versus fluoxetine. A certain drug is most often chosen because of its known properties in the treatment of associated disorders, or BPD symptoms that are also known to be present in other conditions, such as depressive, psychotic, or anxious disorders. However, there are no drugs available for BPD treatment specifically. Available data of the studies included here suggested adverse effects included weight gain, sedation and change of haemogram parameters with olanzapine treatment, and weight loss with topiramate. All drugs were well tolerated in terms of attrition. The data also indicated that there may be an increase in self-harming behaviour in patients treated with olanzapine. Drugs are widely used in borderline personality disorder BPD treatment, chosen because of properties known from other psychiatric disorders "off-label use" , mostly targeting affective or impulsive symptom clusters. Psychological therapies for borderline personality disorder Crisis interventions for people with borderline personality disorder Omega-3 fatty acids for autism spectrum disorders ASD Polyunsaturated fatty acids PUFA supplements for attention deficit hyperactivity disorder ADHD in children and adolescents Polyunsaturated fatty acids PUFAs for children with specific learning disorders. However, these are mostly based on single study effect estimates. Reviews and ratings for olanzapine when used in the treatment of borderline personality disorder. 19 reviews submitted. Oct 1, - In , Marsha Linehan, the creator of Dialectical Behavioral Therapy (DBT), published an article with several colleagues titled: Olanzapine Plus Dialectical Behavior Therapy for Women With High Irritability Who Meet Criteria for Borderline Personality Disorder: A Double-Blind, Placebo-Controlled Pilot. Sep 7, - Olanzapine (Zyprexa) has the most positive data of any medication tested for BPD. The authors review five placebo-controlled trials, all of which showed benefits of olanzapine for aggression, impulsivity, and, though not as consistently, depression. Patients on olanzapine had a mean weight gain of 3 kg. Jun 16, - A certain drug is most often chosen because of its known properties in the treatment of associated disorders, or BPD symptoms that are also known to be adverse effects included weight gain, sedation and change of haemogram parameters with olanzapine treatment, and weight loss with topiramate. The primary objective of this proposed study will be to compare the baseline PET scan to the endpoint scan in 15 BPD patients who have been treated with olanzapine. The comparison of the scans will be done through a statistical image analysis, using a pixel-by-pixel group mean subtraction strategy with appropriate. A dose comparison of olanzapine for the treatment of borderline personality disorder: A week randomized double-blind placebo-controlled study. M.C. Zanarini. x. M.C. Zanarini. Search for articles by this author. My experience taking Zyprexa for bipolar disorder, borderline personality disorder and anxiety. I talk about. Not recommended for the routine use for the treatment of Borderline Personality Disorder. These agents have potential but evidence from pragmatic trials among people using secondary mental health services is lacking and data on long term clinical and cost effectiveness does not exist. N. Olanzapine. If using olanzapine a. Jul 24, - Background Asenapine is a new second-generation antipsychotic that is understudied in borderline personality disorder (BPD). Only one study investigating the use of the drug in this indication (an. Nov 15, - Although many of the medications tested have been demonstrated to be useful, no clear main treatment for BPD has emerged. Despite the efficacy of some of the medicines, acceptability and side effects have proven to be barriers to the

olanzapine borderline personality

use of medication. Therefore, an open-label olanzapine trial utilizing.