

dulcolax suppository generic name

[\[PDF\] is lexapro available as a generic](#)

[\[PDF\] generic brand of depo provera](#)

[\[PDF\] is there a generic for rizatriptan](#)

[\[PDF\] amlodipine price in the philippines](#)

[\[PDF\] metformin 1000 mg cost](#)

[\[PDF\] online rezept f¼r levitra](#)

[\[PDF\] depakote for major depressive disorder](#)

You may also report side effects at [rubeninorchids.com](http://Drug information provided by: Dulcolax - I threw up 20 minutes after taking 2 pills. The information at Drugs. The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Do not use this medicine without a doctor's advice if you are pregnant. This content does not have an English version. Overuse of laxatives can cause your bowels to function improperly, or can make you dependent on laxative use. There are several different types of rectal laxatives and they work in different ways. You should have a bowel movement within 15 to 60 minutes after using a rectal suppository, or within 5 to 20 minutes after using the rectal enema. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Bisacodyl - Is Bisacodyl a substitution for dulcolax? Subscribe to receive email notifications whenever new articles are published. We comply with the HONcode standard for trustworthy health information - verify here.</u>
<u>DULCOLAX SUPPOSITORIES (Bisacodyl) drug information & product resources from MPR including dosage information, educational materials, & patient assistance. Detailed dosage guidelines and administration information for Dulcolax Suppository (bisacodyl). Includes dose adjustments, warnings and precautions. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses. Many times Dulcolax (bisacodyl suppositories) is taken on an as needed basis. Do not take more often than told by the doctor. See also: Dosage Information (in more detail). Dulcolax Suppository official prescribing information for healthcare professionals. Includes: indications, dosage, adverse reactions, pharmacology and more. Physician reviewed Dulcolax (oral and rectal) patient information - includes Dulcolax description, dosage and directions. You should have a bowel movement within 15 to 60 minutes after using a rectal suppository, or within 5 to 20 minutes after using the rectal enema. Do not take a rectal suppository or enema by mouth. DULCOLAX 10 mg Suppositories - Summary of Product Characteristics (SmPC) by SANOFI. Smooth white torpedo shaped suppositories, each containing 10 mg of bisacodyl. 3. In the management of constipation, once regularity has been restarted dosage should be reduced and can usually be stopped. Children aged. Normally the dosage is 5 or 10 mg, but up to 30 mg can be taken for complete cleansing of the bowel before a procedure. When administered rectally in suppository form, it is usually effective in 15 to 60 minutes. Two suppositories can be inserted at once if a very strong, purgative, enema-like result is needed. A few hours. Jan 19, - Dulcolax suppositories melt in the rectum, releasing the bisacodyl to produce its effect on the wall of the colon and rectum. The suppositories start to constipation relief tablets. Bisacodyl gastro-resistant tablets, suppositories and enema are also available without a brand name, ie as the generic medicine. Dulcolax Rectal suppository drug summary. Find medication information including related drug classes, side effects, patient statistics and answers to frequently asked questions. Visit <a href=) for more details. Aug 24, - Bisacodyl should only be used to provide short-term relief from constipation. Suppositories will take between minutes to work, but usually have their effect within 30 minutes. Bisacodyl is sometimes used to clear the bowel before a medical examination which requires the bowel to be empty. When.