

phenergan prescription dosage

[\[PDF\] lasix generic picture](#)

[\[PDF\] lidocaine for sale philippines](#)

[\[PDF\] generic alli orlistat](#)

[\[PDF\] sildenafil free shipping](#)

[\[PDF\] cyclophosphamide cost in india](#)

[\[PDF\] can you take generic tylenol during pregnancy](#)

[\[PDF\] valsartan ratiopharm nebenwirkungen](#)

Be careful if you drive or do anything that requires you to be awake and alert. Available for Android and iOS devices. Phenergan also prevents motion sickness, and treats nausea and vomiting or pain after surgery. Do not take this medicine in larger or smaller amounts or for longer than recommended. Overdose symptoms may include overactive reflexes, loss of coordination, severe drowsiness or weakness, fainting, dilated pupils, weak or shallow breathing, or seizure convulsions. Phenergan should not be given to a child younger than 2 years old. Oral, rectal, IM or IV: Promethazine also acts as an antihistamine. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Subscribe to receive email notifications whenever new articles are published. Phenergan doses are based on weight in children, and any changes may affect your child's dose. Maximum daily dose should not exceed 75 mg. If your dose is different, do not change it unless your doctor tells you to do so. Single mg doses at bedtime or to mg taken three times daily will usually suffice. After initiation of treatment in children or adults, dosage should be adjusted to the smallest amount adequate to relieve symptoms. The administration of promethazine HCl in mg doses will control minor transfusion reactions of an. Allergic Conditions. PO/PR: 25 mg at bedtime or mg before meals and at bedtime (dosage range, mg q8hr). IV/IM: 25 mg; may be repeated in 2 hours when necessary; switch to PO as soon as possible. Nausea & Vomiting. PO/PR: mg qhr PRN. IV/IM: mg qhr PRN. Motion Sickness. allergic conditions. [mg PO tid-qid prn]: Alt: mg IM/IV tid-qid prn; Max: mg/day; Info: give before meals and bedtime; may give second IM/IV dose 2h after first dose; Info: IM preferred over IV. Follow all directions on your prescription label. Your doctor may occasionally change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended. Phenergan is often taken at bedtime or before meals. For motion sickness, Phenergan is usually ?Side Effects ?Dosage ?Phenergan (Promethazine ?Drug Interactions. Do not take Phenergan if you have jaundice. Do not take this medicine if you are receiving or have received high doses of other CNS depressant medications (for example, drugs that are used to treat anxiety and sleep disorders). Any person who is unconscious or in a coma must not be treated with Phenergan. Do not give. The text only version may be available from RNIB in large print, Braille or audio CD. For further information call RNIB Medicine Leaflet Line on The product code(s) for this leaflet is: PL / Phenergan 25 mg tablets. PACKAGE LEAFLET: INFORMATION FOR THE USER. Phenergan 25 mg Tablets. Phenergan Oral tablet mg Drug Medication Dosage information. Learn about the reported side effects, related class drugs, and how these medications will affect your daily lifestyle. Visit rubeninorchids.com for more details. Nov 27, - DESCRIPTION. Promethazine hydrochloride is a white or faintly yellow, practically odourless, crystalline powder. It is very soluble in water, freely soluble in alcohol and in chloroform, and practically insoluble in ether. Phenergan Elixir contains promethazine hydrochloride 5 mg / 5 mL. Phenergan Elixir also. Promethazine Oral tablet 50mg Drug Medication Dosage information. Learn about the reported side effects, related class drugs, and how these medications will affect your daily lifestyle. Visit rubeninorchids.com for more details. Oct 22, - Phenergan is a medicine containing promethazine hydrochloride, an antihistamines used to treat allergy, stop vomiting and for sedation. Dose advice. Dosage varies according to the condition being treated and the individual response to the rubeninorchids.com Disorders: Adults: mg as a single dose at.